

MRI Evaluation of Cartilage Maturation

Miika T. Nieminen, PhD

Department of Diagnostic Radiology, Oulu University Hospital,
Oulu, Finland

miika.nieminen@oulu.fi <http://www.finncartilage.org>

1. Introduction

Various quantitative MRI (qMRI) techniques hold potential in evaluating the extracellular matrix of articular cartilage. In order to apply these techniques in evaluating tissue maturation and to correctly interpret the results it is necessary to understand the structural and compositional changes that occur within growing cartilage. Also, understanding the MRI appearance of cartilage, as verified by histological correlation of tissue content, may help in preventing the pitfalls underlying in the frequent use of immature animal cartilage as a model for human articular cartilage.

This paper summarizes the current understanding of cartilage remodeling during maturation, and the available MRI techniques for its evaluation are reviewed. Selected MRI results of cartilage development are presented with parallel histology in order to interpret the MRI findings.

2. Mature articular cartilage

Adult cartilage is composed of water (60-80% by weight), relatively few chondrocytes (1%), and hydrophilic proteoglycan aggregates (4-7%) enforced by a network of collagen fibrils (15-22%) (1). The proteoglycan concentration increases from surface to deep cartilage (2) while collagen concentration behaves in an inverse manner (3). Benninghoff initially reported the complex structure of the collagen network with three distinct zones from the articular surface to deep tissue: the tangential zone (5-15% of total cartilage thickness) has collagen fibrils aligned along the articular surface, arching into the transitional zone (1-15%) where fibrils have a more or less random orientation, and finally arrange to become perpendicular to the subchondral bone in the radial zone (70-90%) where fibrils become anchored to the underlying bone (4).

To accommodate the varying loading conditions, topographical variation of structure and composition of cartilage exists, resulting in different mechanical properties in different areas of the joint (5). Thus, the ultimate goal in the development of MRI techniques is not only to estimate the construction of the extracellular matrix but to predict the function of the tissue in order to evaluate its prerequisites to accomplish the demanding task of energy absorption and dissipation during locomotion.

3. Immature articular cartilage

While the structure and composition of adult articular cartilage has been known for several decades, only recent advances in histological techniques have provided insight into the changes taking place in the architecture of maturing cartilage. Enhanced polarized light microscopy (ePLM) has recently provided means to study the fiber angle and anisotropy (parallelism) of collagen fibrils (6,7). Fourier Transform Infrared Imaging Spectroscopy (FT-IRIS), on the other hand, enables the assessment of the spatial variation of collagen and proteoglycan contents (8). More conventional techniques, namely the densitometry of stained proteoglycans (9) and biochemical assays (10,11) supplement the aforementioned methods in revealing tissue changes in the course of maturation.

In the developing joint, bone ends are initially formed of epiphyseal hyaline cartilage, and become ossified with age apart from the layer of articular cartilage. Unlike adult articular cartilage, postnatal human cartilage is lacking an organized collagen network. Collagen fibrils of young porcine cartilage (4 months) are reported to exhibit a preferential tangential arrangement even in deep parts of the tissue, yet the order of anisotropy in the radial zone is lower than in adult cartilage (12). In immature tissue, several additional histological zones (up to seven in total) have been observed in juvenile animal tissue (7,13-16), and these are possibly related to zones of high cell density where collagen fibrils curve around the chondrocytes (7,16). These tissue areas most likely represent the cartilaginous epiphysis which will eventually become bone. Both the spatial distribution and content of collagen modulates during the development of cartilage, reaching its highest content at maturity (12). On the contrary, a significant decrease in proteoglycan content has been reported with maturation in rat cartilage (17). The overall solid content slowly increases with maturation (i.e. water content decreases) (18). The significant increase of confined compression modulus and the decrease in hydraulic permeability of fetal to adult bovine cartilage are primarily associated to the development of the collagen component (18). Thus, during skeletal maturation towards the final cartilage phenotype

the tissue structure and composition go through phases of significant remodeling that are dramatically reflected in the functional properties of the tissue.

4. qMRI for cartilage composition and structure

Several qMRI techniques have been tested in assessing the collagen component of cartilage. T_2 relaxation time was initially the parameter of interest due to the distinct and unique MRI appearance of cartilage on T_2 -weighted MR images (19). Cartilage T_2 is sensitive to the architecture and organization of collagen fibrils (16,20,21), but also dependent on the collagen content (22,23) and water content of the tissue (24,25). Conflicting reports exist on the influence of cartilage proteoglycans on T_2 relaxation (26,27). While T_2 relaxation time may not be specific to collagen, its spatial variation and laminar appearance is indisputably dependent on the arrangement of the collagen fibrils in the B_0 field due to the orientation dependent dipolar interaction of collagen associated water, known as the “magic angle effect”. The in vivo application of T_2 is challenged by the varying orientation of the curved articular surfaces that lead to an orientation dependent variation of T_2 even in normal cartilage. Yet, promising in vivo results have been reported in the recent years (28). In addition to T_2 relaxation, also other MRI parameters have shown a degree of sensitivity on the collagen component of cartilage. The collagen matrix is the predominant determinant of magnetization transfer effects in cartilage, however, it is not very sensitive to changes in the physiological range (29,30). Diffusion tensor imaging of cartilage may enable the determination of the collagen fibril arrangement (31). Similar to T_2 , ^2H and ^{23}Na quadrupolar splittings depend on the direction of interaction and the B_0 field and can be used to study the orientation of fibrils in different cartilage zones (32).

For the estimation of the proteoglycan content of cartilage, the delayed Gadolinium Enhanced MRI of Cartilage (dGEMRIC) technique has been widely used. The technique utilizes the cationic contrast agent Gd-DTPA^{2-} which distributes in inverse relation to the fixed negative charge of the glycosaminoglycans of proteoglycans in order to modulate the T_1 relaxation time of the tissue (33). The technique has been validated both for bulk tissue and spatially (34,35), and has been applied in several in vitro and in vivo studies (36-45). $T_{1\rho}$ mapping, i.e. T_1 relaxation in the rotating frame, is another candidate for evaluating cartilage proteoglycans (46). It shows a high correlation with biochemical proteoglycan measures (47,48) and initial in vivo testing has been reported (49,50). The specificity of $T_{1\rho}$ for cartilage proteoglycans, however, is hindered by its dependence on residual dipolar interaction related to collagen orientation (51). Finally, Na-MRI is a method that can be used to quantify the amount of sodium ions attracted by the fixed negative charge of cartilage proteoglycans (52). The technique is sensitive and specific in detecting small changes in proteoglycan concentration (53) and has been tested in an in vivo model (54).

Mechanical properties of cartilage have been successfully related to dGEMRIC, T_2 and $T_{1\rho}$ (38,39,43,55-58). The correlations between moduli and MRI parameters has been high at best, and the techniques have reproduced the topographical variation in mechanical properties across the joint (39,59). The correlations, however, tend to be inferior when examining pathologic tissue.

5. qMRI of the developing extracellular matrix

In the wake of microscopic discoveries of cartilage maturation, MRI has shown its potential in assessing cartilage development. T_2 -weighted signal is reported to decrease with maturation, and the homogeneous appearance of the cartilaginous epiphysis at the very early childhood is reported to change to a laminated appearance resembling that of mature cartilage (60,61). With reference to the findings presented in section 3, the signal decrease is likely caused by the remodeling of the collagen network towards an anisotropic structure and by the general increase in solid content (collagen and proteoglycans).

T_2 relaxation time mapping has shown promise in revealing the diverging cartilage structure. The additional zones observed with polarized light microscopy correlate well with the depth-wise variation of T_2 relaxation time (7,13,16). These structures frequently exhibit consecutive zones of low and high collagen anisotropy, and they are manifested in MRI as laminae of long and short T_2 relaxation times, respectively (Fig. 1A). The probable mechanisms for the T_2 elevation is the reduced dipolar interaction due to lower collagen anisotropy and increased water content due to hypercellularity (16). Complicated collagenous structures in the young canine humeral head has been revealed: fibrils in central areas exhibit a tri-laminar appearance while the structure becomes more complicated when moving to the peripheral area of the joint with different loading conditions (7). This could possibly explain the higher T_2 relaxation time observed in non-weight-bearing cartilage as compared to weight-bearing tissue in the femoral compartment of the pediatric knee (62). A significant increase in ^2H and ^{23}Na quadrupolar splitting with age has been observed, providing another means to study the order of collagen anisotropy with maturation (63).

dGEMRIC has been used to investigate the proteoglycan content in the course of cartilage development in a single study (64). Histology of superficial cartilage in adult humans showed a lower proteoglycan content as

opposed to immature bovine and immature porcine cartilage. dGEMRIC revealed a similar trend, however, the difference was not statistically significant. A poor correlation between dGEMRIC and histological quantitation of stained proteoglycans has been revealed in the deep parts of immature cartilage (35,64). This is possibly related to the high cellularity of deep juvenile cartilage which infringes the single compartment assumption made in the dGEMRIC technique or high tissue permeability in these tissue zones that could restrict the accumulation of the contrast agent (35). Nonetheless, several other studies have shown the capability of dGEMRIC in revealing subtle changes in the PG content of cartilage (Fig. 1B), and when used with caution it is one of the most potential techniques in revealing maturation related proteoglycan changes in cartilage.




Figure 1: For a sample of juvenile cartilage imaged at 9.4T (cartilage surface is towards the top of the page), **A:** T_2 map and ePLM images (orientation, anisotropy and birefringence); **zone I:** superficial cartilage with collagen fibrils arranged along the cartilage surface, and a concomitant short T_2 relaxation time; **zone II:** arcading collagen fibrils with low anisotropy together with a long T_2 ; **zone III:** anisotropic collagen fibrils with tangential arrangement (as in zone I) and short T_2 relaxation time; **zone IV:** anisotropic collagen arrangement and prolonged T_2 ; **zone V:** radial collagen fibrils with some degree of anisotropy and short T_2 relaxation time. **B:** safranin-O stained section and respective dGEMRIC map: proteoglycan content increases from superficial to deep cartilage and this variation is also visualized by dGEMRIC. (Image courtesy of Mikko Nissi, University of Kuopio)

Surgically produced repair tissue, such as that resulting from autologous chondrocyte transplantation (ACT), is analogous to postnatal articular cartilage in the sense that it lacks the three-dimensional arrangement of collagen fibrils. dGEMRIC and T_2 mapping have been used to evaluate ACT repair tissue in humans (36,37,65). dGEMRIC revealed PG replenishment in ACT grafts. The repair tissue, however, showed an elevated T_2 relaxation time without a distinct laminar appearance, which may be explained by the lack of the three-dimensional collagenous architecture. T_2 mapping has the potential to show emerging arrangement of collagen in repair tissue, however, it is unlikely that the tissue develops such a structure any quicker than in the age of skeletal maturation. Therefore long term follow-up qMRI studies are required.

Another study used various MRI parameters to evaluate regenerated tissue in an animal model of spontaneous repair (66). Repair tissue showed a trend toward low PG content, as assessed by dGEMRIC, and low T_2 values likely resulting from the fibrous cartilage structure as verified by polarized light microscopy. T_1 and diffusion measurements were not particularly sensitive in differentiating regenerated tissue from normal cartilage, however, the number of samples in the study was very small and prevented from running statistical tests.

6. Summary

Developing cartilage shows dramatic changes in the extracellular matrix in course of time. qMRI techniques are very promising in the nondestructive assessment of the maturation related changes in the macromolecular constituents of cartilage. To date, T_2 mapping and dGEMRIC have been used in revealing structural

changes in the three-dimensional collagen matrix and proteoglycans, respectively. Several other MRI parameters have the potential to further characterize cartilage maturation and their applicability remains to be shown.

Acknowledgments

Jarno Rieppo, MB, and Mika Hyttinen, MD, are acknowledged for fruitful discussion and sharing their latest results, and the members of the FinnCartilage consortium for productive collaboration.

References

1. Mow VC, Ratcliffe A, Poole AR. Cartilage and diarthrodial joints as paradigms for hierarchical materials and structures. *Biomaterials* 1992;13(2):67-97.
2. Maroudas A, Bayliss MT, Venn MF. Further studies on the composition of human femoral head cartilage. *Ann Rheum Dis* 1980;39(5):514-523.
3. Venn M, Maroudas A. Chemical composition and swelling of normal and osteoarthrotic femoral head cartilage. I. Chemical composition. *Ann Rheum Dis* 1977;36(2):121-129.
4. Benninghoff A. Form und bau der Gelenkknorpel in ihren Beziehungen zur Function. Erste Mitteilung: die modellierenden und formenhaltenden Faktoren des Knorpelreliefs. *Z Anat* 1925;76:43-63.
5. Jurvelin JS, Arokoski JP, Hunziker EB, Helminen HJ. Topographical variation of the elastic properties of articular cartilage in the canine knee. *J Biomech* 2000;33(6):669-675.
6. Rieppo J, Hallikainen J, Jurvelin JS, Helminen HJ, Hyttinen MM. Novel quantitative polarization microscopic assessment of cartilage and bone collagen birefringence, orientation and anisotropy. *Transact Orthop Res Soc* 2003;28:570.
7. Xia Y, Moody JB, Alhadlaq H, Hu J. Imaging the physical and morphological properties of a multi-zone young articular cartilage at microscopic resolution. *J Magn Reson Imaging* 2003;17(3):365-374.
8. Camacho NP, West P, Torzilli PA, Mendelsohn R. FTIR microscopic imaging of collagen and proteoglycan in bovine cartilage. *Biopolymers* 2001;62(1):1-8.
9. Kiviranta I, Jurvelin J, Tammi M, Säämänen AM, Helminen HJ. Microspectrophotometric quantitation of glycosaminoglycans in articular cartilage sections stained with Safranin O. *Histochemistry* 1985;82(3):249-255.
10. Goldberg RL, Kolibas LM. An improved method for determining proteoglycans synthesized by chondrocytes in culture. *Connect Tissue Res* 1990;24(3-4):265-275.
11. Woessner JF. The determination in tissue and protein samples containing small proportions of this imino acid. *Arch Biochem Biophys* 1961;93:440-447.
12. Rieppo J, Hyttinen MM, Halmesmäki E, Ruotsalainen H, Vasara A, Kiviranta I, Jurvelin JS, Helminen HJ. Remodelation of Collagen Network Architecture during Cartilage Maturation. *Transact Orthop Res Soc* 2004;29:549.
13. Nissi MJ, Rieppo J, Töyräs J, Laasanen MS, Helminen HJ, Jurvelin JS, Nieminen MT. T₂ relaxation reveals differences in spatial collagen network anisotropy in human, bovine and porcine articular cartilage. *Proc Intl Soc Magn Reson Med* 2003;11:54.
14. Xia Y, Moody JB, Alhadlaq H, Burton-Wurster N, Lust G. Characteristics of topographical heterogeneity of articular cartilage over the joint surface of a humeral head. *Osteoarthritis Cartilage* 2002;10(5):370-380.
15. Gründer W, Kanowski M, Wagner M, Werner A. Visualization of pressure distribution within loaded joint cartilage by application of angle-sensitive NMR microscopy. *Magn Reson Med* 2000;43(6):884-891.
16. Nieminen MT, Töyräs J, Rieppo J, Silvennoinen MJ, Hakumäki JM, Hyttinen MM, Helminen HJ, Jurvelin JS. T₂ relaxation reveals spatial collagen architecture in articular cartilage: a comparative quantitative MRI and polarized light microscopic study. *Magn Reson Med* 2001;46:487-493.
17. Watrin A, Ruaud JP, Olivier PT, Guingamp NC, Gonord PD, Netter PA, Blum AG, Guillot GM, Gillet PM, Loeuille DH. T₂ mapping of rat patellar cartilage. *Radiology* 2001;219(2):395-402.
18. Williamson AK, Chen AC, Sah RL. Compressive properties and function-composition relationships of developing bovine articular cartilage. *J Orthop Res* 2001;19(6):1113-1121.
19. Rubenstein JD, Kim JK, Morova-Protzner I, Stanchev PL, Henkelman RM. Effects of collagen orientation on MR imaging characteristics of bovine articular cartilage. *Radiology* 1993;188(1):219-226.
20. Gründer W, Wagner M, Werner A. MR-microscopic visualization of anisotropic internal cartilage structures using the magic angle technique. *Magn Reson Med* 1998;39(3):376-382.
21. Xia Y, Moody JB, Burton-Wurster N, Lust G. Quantitative in situ correlation between microscopic MRI and polarized light microscopy studies of articular cartilage. *Osteoarthritis Cartilage* 2001;9(5):393-406.
22. Fragonas E, Mlynarik V, Jellus V, Micali F, Piras A, Toffanin R, Rizzo R, Vittur F. Correlation between biochemical composition and magnetic resonance appearance of articular cartilage. *Osteoarthritis Cartilage* 1998;6(1):24-32.
23. Menezes NM, Gray ML, Hartke JR, Burstein D. T₂ and T1rho MRI in articular cartilage systems. *Magn Reson Med* in press.
24. Lusse S, Claassen H, Gehrke T, Hassenpflug J, Schunke M, Heller M, Gluier CC. Evaluation of water content by spatially resolved transverse relaxation times of human articular cartilage. *Magnetic Resonance Imaging* 2000;18(4):423-430.
25. Shapiro EM, Borthakur A, Kaufman JH, Leigh JS, Reddy R. Water distribution patterns inside bovine articular cartilage as visualized by 1H magnetic resonance imaging. *Osteoarthritis Cartilage* 2001;9(6):533-538.
26. Nieminen MT, Töyräs J, Rieppo J, Hakumäki JM, Silvennoinen J, Helminen HJ, Jurvelin JS. Quantitative MR microscopy of enzymatically degraded articular cartilage. *Magn Reson Med* 2000;43(5):676-681.

27. Wayne JS, Kraft KA, Shields KJ, Yin C, Owen JR, Disler DG. MR imaging of normal and matrix-depleted cartilage: correlation with biomechanical function and biochemical composition. *Radiology* 2003;228(2):493-499.
28. Mosher TJ, Dardzinski BJ. Cartilage MRI T₂ relaxation time mapping: overview and applications. *Semin Musculoskelet Radiol* 2004;8(4):355-368.
29. Kim DK, Ceckler TL, Hascall VC, Calabro A, Balaban RS. Analysis of water-macromolecule proton magnetization transfer in articular cartilage. *Magn Reson Med* 1993;29(2):211-215.
30. Gray ML, Burstein D, Lesperance LM, Gehrke L. Magnetization transfer in cartilage and its constituent macromolecules. *Magn Reson Med* 1995;34(3):319-325.
31. Filidoro L, Dietrich O, Weber J, Rauch E, Oerther T, Wick M, Reiser MF, Glaser C. High-resolution diffusion tensor imaging of human patellar cartilage: feasibility and preliminary findings. *Magn Reson Med* 2005;53(5):993-998.
32. Navon G, Shinar H, Eliav U, Seo Y. Multiquantum filters and order in tissues. *NMR Biomed* 2001;14(2):112-132.
33. Bashir A, Gray ML, Burstein D. Gd-DTPA²⁻ as a measure of cartilage degradation. *Magn Reson Med* 1996;36(5):665-673.
34. Bashir A, Gray ML, Hartke J, Burstein D. Nondestructive imaging of human cartilage glycosaminoglycan concentration by MRI. *Magn Reson Med* 1999;41(5):857-865.
35. Nieminen MT, Rieppo J, Silvennoinen J, Töyräs J, Hakumäki JM, Hyttinen MM, Helminen HJ, Jurvelin JS. Spatial assessment of articular cartilage proteoglycans with Gd-DTPA²⁻ enhanced T₁ imaging. *Magn Reson Med* 2002;48(4):640-648.
36. Gillis A, Bashir A, McKeon B, Scheller A, Gray ML, Burstein D. Magnetic resonance imaging of relative glycosaminoglycan distribution in patients with autologous chondrocyte transplants. *Invest Radiol* 2001;36(12):743-748.
37. Vasara AI, Nieminen MT, Jurvelin JS, Peterson L, Lindahl A, Kiviranta I. Indentation stiffness of repair tissue after autologous chondrocyte transplantation. *Clin Orthop Relat Res* 2005(433):233-242.
38. Nissi MJ, Töyräs J, Laasanen MS, Rieppo J, Saarakkala S, Lappalainen R, Jurvelin JS, Nieminen MT. Proteoglycan and collagen sensitive MRI evaluation of normal and degenerated articular cartilage. *J Orthop Res* 2004;23:557-564.
39. Kurkijärvi JE, Nissi MJ, Kiviranta I, Jurvelin JS, Nieminen MT. Delayed gadolinium-enhanced MRI of cartilage (dGEMRIC) and T₂ characteristics of human knee articular cartilage: topographical variation and relationships to mechanical properties. *Magn Reson Med* 2004;52:41-46.
40. Tiderius CJ, Olsson LE, Leander P, Ekberg O, Dahlberg L. Delayed gadolinium-enhanced MRI of cartilage (dGEMRIC) in early knee osteoarthritis. *Magn Reson Med* 2003;49(3):488-492.
41. Tiderius CJ, Svensson J, Leander P, Ola T, Dahlberg L. dGEMRIC (delayed gadolinium-enhanced MRI of cartilage) indicates adaptive capacity of human knee cartilage. *Magn Reson Med* 2004;51(2):286-290.
42. Kim YJ, Jaramillo D, Millis MB, Gray ML, Burstein D. Assessment of early osteoarthritis in hip dysplasia with delayed gadolinium-enhanced magnetic resonance imaging of cartilage. *J Bone Joint Surg Am* 2003;85-A(10):1987-1992.
43. Lammintausta E, Kiviranta P, Nissi MJ, Laasanen MS, Kiviranta I, Nieminen MT, Jurvelin JS. T₂ Relaxation Time and delayed Gadolinium-Enhanced MRI of cartilage (dGEMRIC) of human patellar cartilage at 1.5 T and 9.4 T: correlation with tissue mechanical properties. *J Orthop Res*, in press.
44. Williams A, Gillis A, McKenzie C, Po B, Sharma L, Micheli L, McKeon B, Burstein D. Glycosaminoglycan distribution in cartilage as determined by delayed gadolinium-enhanced MRI of cartilage (dGEMRIC): potential clinical applications. *AJR Am J Roentgenol* 2004;182(1):167-172.
45. Williams A, Sharma L, McKenzie CA, Prasad PV, Burstein D. Delayed gadolinium-enhanced magnetic resonance imaging of cartilage in knee osteoarthritis: Findings at different radiographic stages of disease and relationship to malalignment. *Arthritis Rheum* 2005;52(11):3528-3535.
46. Duvvuri U, Reddy R, Patel SD, Kaufman JH, Kneeland JB, Leigh JS. T1rho-relaxation in articular cartilage: effects of enzymatic degradation. *Magn Reson Med* 1997;38(6):863-867.
47. Akella SV, Regatte RR, Gougoutas AJ, Borthakur A, Shapiro EM, Kneeland JB, Leigh JS, Reddy R. Proteoglycan-induced changes in T1rho-relaxation of articular cartilage at 4T. *Magn Reson Med* 2001;46(3):419-423.
48. Duvvuri U, Kudchodkar S, Reddy R, Leigh JS. T(1rho) relaxation can assess longitudinal proteoglycan loss from articular cartilage in vitro. *Osteoarthritis Cartilage* 2002;10(11):838-844.
49. Regatte RR, Akella SV, Wheaton AJ, Lech G, Borthakur A, Kneeland JB, Reddy R. 3D-T1rho-relaxation mapping of articular cartilage: in vivo assessment of early degenerative changes in symptomatic osteoarthritic subjects. *Acad Radiol* 2004;11(7):741-749.
50. Regatte RR, Akella SV, Borthakur A, Kneeland JB, Reddy R. In vivo proton MR three-dimensional T1rho mapping of human articular cartilage: initial experience. *Radiology* 2003;229(1):269-274.
51. Akella SV, Regatte RR, Wheaton AJ, Borthakur A, Reddy R. Reduction of residual dipolar interaction in cartilage by spin-lock technique. *Magn Reson Med* 2004;52(5):1103-1109.
52. Lesperance LM, Gray ML, Burstein D. Determination of fixed charge density in cartilage using nuclear magnetic resonance. *J Orthop Res* 1992;10(1):1-13.
53. Borthakur A, Shapiro EM, Beers J, Kudchodkar S, Kneeland JB, Reddy R. Sensitivity of MRI to proteoglycan depletion in cartilage: comparison of sodium and proton MRI. *Osteoarthritis Cartilage* 2000;8(4):288-293.
54. Wheaton AJ, Borthakur A, Dodge GR, Kneeland JB, Schumacher HR, Reddy R. Sodium magnetic resonance imaging of proteoglycan depletion in an in vivo model of osteoarthritis. *Acad Radiol* 2004;11(1):21-28.
55. Nieminen MT, Töyräs J, Laasanen MS, Silvennoinen J, Helminen HJ, Jurvelin JS. Prediction of biomechanical properties of articular cartilage with quantitative magnetic resonance imaging. *J Biomech* 2004;37(3):321-328.

56. Samosky JT, Burstein D, Eric Grimson W, Howe R, Martin S, Gray ML. Spatially-localized correlation of dGEMRIC-measured GAG distribution and mechanical stiffness in the human tibial plateau. *J Orthop Res* 2005;23(1):93-101.
57. Wheaton AJ, Dodge GR, Elliott DM, Nicoll SB, Reddy R. Quantification of cartilage biomechanical and biochemical properties via T(1rho) magnetic resonance imaging. *Magn Reson Med* 2005;54(5):1087-1093.
58. Wayne JS, Kraft KA, Shields KJ, Yin C, Owen JR, Disler DG. MR imaging of normal and matrix-depleted cartilage: correlation with biomechanical function and biochemical composition. *Radiology* 2003;228(2):493-499.
59. Nieminen MT, Töyräs J, Laasanen MS, Silvennoinen J, Helminen HJ, Jurvelin JS. Prediction of biomechanical properties of articular cartilage with quantitative magnetic resonance imaging. *J Biomech* 2004;37(3):321-328.
60. Varich LJ, Laor T, Jaramillo D. Normal maturation of the distal femoral epiphyseal cartilage: age-related changes at MR imaging. *Radiology* 2000;214(3):705-709.
61. Babyn PS, Kim HK, Lemaire C, Gahunia HK, Cross A, DeNanassy J, Pritzker KP. High-resolution magnetic resonance imaging of normal porcine cartilaginous epiphyseal maturation. *J Magn Reson Imaging* 1996;6(1):172-179.
62. Dardzinski BJ, Laor T, Schmithorst VJ, Klosterman L, Graham TB. Mapping T₂ relaxation time in the pediatric knee: feasibility with a clinical 1.5-T MR imaging system. *Radiology* 2002;225(1):233-239.
63. Keinan-Adamsky K, Shinar H, Navon G. A Multinuclear NMR Study of the Maturation of Pig Articular Cartilage. *Proc Intl Soc Magn Reson Med* 2003;11:1503.
64. Nissi MJ, Rieppo J, Töyräs J, Laasanen MS, Kiviranta I, Jurvelin JS, Nieminen MT. MRI quantitation of proteoglycans with dGEMRIC in human, bovine and porcine articular cartilage. *Transact Orthop Res Soc* 2004;29:1004.
65. Kurkijärvi JE, Nissi M, Ojala RO, Vasara AI, Jurvelin JS, Kiviranta I, Nieminen MT. In Vivo T₂ Mapping and dGEMRIC of Human Articular Cartilage Repair after Autologous Chondrocyte Transplantation. *Proc Intl Soc Magn Reson Med* 2005;13:481.
66. Nieminen MT, Winalski CS, Rieppo J, Lammi M, Lechpammer S, Johnson K, Minas T, Kiviranta I. Multi-parametric MRI assessment of cartilage repair with correlation to histology. *Proc Intl Soc Magn Reson Med* 2004;11:2399.